

The problems of determining the Breit–Wigner parameters of nucleon resonance $S_{11}(1535)$

E.V.Balandina, E.M.Leikin and N.P.Yudin

Scobeltsyn Institute of Nuclear Physics,

Moscow State University,

Russia

1 Introduction

This paper completes the phenomenological analysis of the set of experimental data on η -meson photoproduction on protons. The main goal of the analysis was to receive the reliable estimates of some fundamental parameters, first of all the multipole amplitudes of the process. The most direct and statistically justified way is to divide the whole analysis into two stages. The first one is the partial waves analysis of the experimentally measured observables, for example, angular distributions. The contribution of different partial waves can be obtained by linear regression procedure based on linear nonparametric model. Expanding the observables into series, in this case, over the orthogonal Legendre polynomials we can determine the terms significantly different from zero and reveal partial waves participating in the process. Thus, the first stage of analysis give us the number of the linear regression coefficients that provide accordingly statistics the best description of the experimental observables.

2 Results of partial wave analysis

The result of the partial wave analysis of the angular distributions measured by three experimental groups [1, 2] and [3] was presented in [4]. The main conclusions from this analysis are:

1. One confirms the known fact that in energy region from threshold to about 1GeV η -mesons are produced mainly in s -state and the process has the resonance character.
2. The contributions of higher partial waves appear only as sp - and sd -interference and their contributions are small compared to s -wave. Unfortunately the results of three experiments in this respect qualitatively contradict with each other.

We made also the partial wave analysis of angular distribution measured in gamma ray region 0.795 – 1.925 GeV [5]. In this case like the lower energy region the linear model with first three terms in the expansion of the differential cross section of the process over Legendre polynomials was proved to be also statistically justified. The energy dependence of corresponding regression coefficients a_0 , a_1 and a_2 is shown of Fig.1 [6]. These figures confirm the dominant role of the s -wave and presence the p - and d -partial waves by order of magnitude smaller. There is no evidence of resonance contribution besides $S_{11}(1535)$, firmly established in energy region between 1 and 2 GeV, and namely $S_{11}(1650)$, $D_{15}(1675)$, $F_{15}(1680)$ and $P_{13}(1720)$ [7].

Smooth energy behaviour of the regression coefficients a_0 , a_1 and a_2 seems to indicate that in this energy region the principal role are playing the nonresonant waves.




Figure 1: The energy dependence of the regression coefficients a_0 , a_1 , a_2 (W – total energy in CM system) from [5]

3 Breit–Wigner characteristics of $S_{11}(1535)$ -resonance and E_{0+} -amplitude of η -meson photoproduction on protons

The results of the partial wave analysis of the photoproduction of η -mesons on protons in the energy region from threshold to about 2 GeV permit the significant simplification of the second part of the phenomenological analysis that is determination of the multipole amplitudes and parameters of $S_{11}(1535)$ resonance. The statistically justified description of experimental observables with three terms of the expansion over Legendre polynomials means s -wave dominance. So in the expression of a_0 , a_1 and a_2 in terms of multipole amplitudes presented in [4] we can keep only the terms with E_{0+} . The main contribution of s -wave contains a_0 :

$$a_0(W) = |E_{0+}|^2, \quad (1)$$

W – total energy in CM-system.

Assuming the Breit–Wigner behaviour of the electric dipole amplitude we can use this fact as the parametric statistical model for description of regression coefficient $a_0(W)$ or the total cross section $(k/q)\sigma(W) = 4\pi a_0(W)$, where k and q – CM-momenta of gamma quantum and η -meson correspondingly. The relevant formulae that take into account the energy dependence of the resonance width are [1]:

$$\frac{k}{q}\sigma(W) = 4\pi a_0(W) = \frac{\sigma(W_R)W_R^2\Gamma_R^2}{(W_R^2 - W^2)^2 + W_R^2\Gamma^2(W)}, \quad (2)$$

$$\Gamma(W) = \Gamma_R(b_\eta \frac{q_\eta}{q_{\eta R}} + b_\pi \frac{q_\pi}{q_{\pi R}} + b_{\pi\pi}). \quad (3)$$

where Γ is total width of the resonance, b_η , b_π , $b_{\pi\pi}$ are the probabilities of decay to corresponding channels, q_η , q_π are the momenta η and π , index R means the value at resonance. There were three fitted parameters: $\sigma(W_R)$, W_R and Γ_R , which were used to calculate also helicity amplitude:

$$A_{1/2}^2 = \frac{W_R}{2m_p} \frac{\Gamma_R}{b_\eta} \sigma(W_R), \quad (4)$$

(m_p – proton mass) and parameter ξ that doesn't depend on b_η

$$\xi = \frac{1}{2} \sqrt{\frac{k}{q} \sigma(W_R)}. \quad (5)$$

The values of b_η , b_π , $b_{\pi\pi}$ were fixed. The results are presented in tables 1-4. Table 1 presents the estimates received by fitting both $4\pi a_0(W)$ and $(k/q)\sigma(W)$ which agree quite well. But there is the large inconsistency between total width of $S_{11}(1535)$ resonance obtained from the experimental data of paper [1] and [2] and experimental data of papers [3] and [5].

It should be noted that the later estimates agree with values of Γ obtained from hadron processes and thereby remove the disagreement existed in values of Γ estimated from photo- and hadroproduction of η -mesons [7]. The differences between total width and other parameters resulting from experimental data of various experiments are illustrating by Breit-Wigner curves fitted with $b_\eta = 0.55$, $b_\pi = 0.35$ on the figures 2-4.

The large values of criterion χ^2/ν per degree of freedom in tables 2 and 4 is due to the considerable deviation of the experimental values relative the fitted curve which significantly exceed the experimental errors. This means that the deviations of points from the Breit-Wigner curve do not follow the normal distribution with zero expectation. In this case the goodness of fit can be estimated by the value of noncentral criterion χ_{nc}^2




Figure 2: The comparison of experimental points from [1] with parametric model (2), (3) with $b_\eta = 0.55$, $b_\pi = 0.35$.




Figure 3: The comparison of experimental points from [3] with parametric model (2), (3) with $b_\eta = 0.55$, $b_\pi = 0.35$.

Table 1: Parameters of resonance $S_{11}(1535)$ from the [1]

$(k/q)(d\sigma/d\Omega)$	$b_\eta = 0.55, b_\pi = 0.35$	$b_\eta = 0.45, b_\pi = 0.45$	$b_\eta = 0.35, b_\pi = 0.55$
$\sigma(W_R), \mu b$	33.51 ± 1.5	35.88 ± 1.48	38.18 ± 1.57
$W_R, \text{ MeV}$	1563.68 ± 9.53	1555.32 ± 8.32	1547.68 ± 7.86
$\Gamma_R, \text{ MeV}$	270.42 ± 41.18	264.02 ± 40.5	271.34 ± 46.61
$\xi, 10^{-4} \text{ MeV}^{-1}$	2.08 ± 0.05	2.15 ± 0.04	2.22 ± 0.05
$A_{1/2}, 10^{-3} \text{ GeV}^{-1/2}$	129.25 ± 6.78	142.32 ± 7.46	164.28 ± 9.53
χ^2/ν	1.2	1.21	1.24
$4\pi a_0(W)$	$b_\eta = 0.55, b_\pi = 0.35$	$b_\eta = 0.45, b_\pi = 0.45$	$b_\eta = 0.35, b_\pi = 0.55$
$\sigma(W_R), \mu b$	34.63 ± 0.82	36.92 ± 0.8	39.25 ± 0.76
$W_R, \text{ MeV}$	1558.73 ± 4.79	1551.32 ± 4.16	1543.9 ± 3.57
$\Gamma_R, \text{ MeV}$	244.09 ± 20.32	238.41 ± 19.83	240.16 ± 20.44
$\xi, 10^{-4} \text{ MeV}^{-1}$	2.11 ± 0.02	2.18 ± 0.02	2.25 ± 0.02
$A_{1/2}, 10^{-3} \text{ GeV}^{-1/2}$	122.96 ± 3.51	135.31 ± 3.83	154.43 ± 4.42
χ^2/ν	0.52	0.52	0.49

Table 2: Parameters of resonance $S_{11}(1535)$ from [2]

$4\pi a_0(W)$	$b_\eta = 0.55, b_\pi = 0.35$	$b_\eta = 0.45, b_\pi = 0.45$	$b_\eta = 0.35, b_\pi = 0.55$
$\sigma(W_R), \mu b$	41.06 ± 1.24	42.89 ± 1.28	44.79 ± 1.35
$W_R, \text{ MeV}$	1537.33 ± 1.47	1533.53 ± 1.54	1529.4 ± 1.69
$\Gamma_R, \text{ MeV}$	139.7 ± 5.76	137.08 ± 5.49	134.99 ± 5.3
$\xi, 10^{-4} \text{ MeV}^{-1}$	2.3 ± 0.03	2.35 ± 0.04	2.4 ± 0.04
$A_{1/2}, 10^{-3} \text{ GeV}^{-1/2}$	93.43 ± 1.54	102.71 ± 1.67	115.62 ± 1.9
χ^2/ν	15.27	15.29	15.54

with noncentrality parameters δ^2 equal to the sum of expectation squared of individual variables [8]. As an example in Table 5 the differences Δ between the values $4\pi a_0(W)$ obtained as a result of partial waves analysis

Table 3: Parameters of resonance $S_{11}(1535)$ from [3]

$4\pi a_0(W)$	$b_\eta = 0.55, b_\pi = 0.35$	$b_\eta = 0.45, b_\pi = 0.45$	$b_\eta = 0.35, b_\pi = 0.55$
$\sigma(W_R), \mu b$	29.83 ± 0.76	31.16 ± 0.71	32.51 ± 0.66
$W_R, \text{ MeV}$	1559.18 ± 2.83	$1554. \pm 2.46$	1548.47 ± 2.13
$\Gamma_R, \text{ MeV}$	199.61 ± 13.86	196.17 ± 13.43	194.25 ± 13.25
$\xi, 10^{-4} \text{ }^{-1}$	1.96 ± 0.02	$2. \pm 0.02$	2.05 ± 0.02
$A_{1/2}, 10^{-3} \text{ GeV}^{-1/2}$	103.35 ± 2.45	113.83 ± 2.63	128.63 ± 2.92
χ^2/ν	2.3	2.3	2.31

 Table 4: Parameters of resonance $S_{11}(1535)$ from [5]

$4\pi a_0(W)$	$b_\eta = 0.55, b_\pi = 0.35$	$b_\eta = 0.45, b_\pi = 0.45$	$b_\eta = 0.35, b_\pi = 0.55$
$\sigma(W_R), \mu b$	43.18 ± 9.13	46.15 ± 11.43	49.25 ± 14.19
$W_R, \text{ MeV}$	1530.63 ± 12.13	1525.6 ± 13.74	1520.34 ± 15.46
$\Gamma_R, \text{ MeV}$	152.4 ± 18.7	147.94 ± 16.4	143.97 ± 14.53
$\xi, 10^{-4} \text{ }^{-1}$	2.36 ± 0.25	2.44 ± 0.3	2.52 ± 0.36
$A_{1/2}, 10^{-3} \text{ GeV}^{-1/2}$	96.86 ± 8.32	106.03 ± 10.47	118.47 ± 13.74
χ^2/ν	11.22	11.22	11.25

of experimental data [2] and the Breit–Wigner curve for the case $b_\eta = 0.55, b_\pi = 0.35$ are shown.

These differences were used to estimate the value of noncentrality parameter $\delta^2 = 46.8$. The expectation of noncentral value χ_{nc}^2 per degree of freedom turns out to be 1.05 supporting statistical reliability of the result presented in tables 2 and 4. At the same time this item may be considered as some indication of possible biases in experimental data.

Description of energy dependence $4\pi a_0(W)$ by parametric model based on the Breit–Wigner formula allows to find the real and imaginary parts




Figure 4: The comparison of experimental points from [2] (a) and [5] (b) with parametric model (2), (3) with $b_\eta = 0.55$, $b_\pi = 0.35$.

Table 5: The differences Δ between experimental and fitted values of $4\pi a_0(W)$ from [2].

W , MeV	Δ	W , MeV	Δ	W , MeV	Δ
1490.16	2.9	1553.9	1.41	1612.92	0.83
1501.14	0.36	1564.08	0.98	1622.26	-0.06
1511.98	0.09	1574.06	0.82	1631.55	-0.67
1522.61	-2.6	1583.98	2.3	1640.78	-1.09
1533.85	-1.59	1593.49	2.78	1649.79	-1.53
1543.61	-1.06	1603.12	2.03	1658.81	-1.55

and also the phase of electric dipole amplitude E_{0+} responsible for production of $S_{11}(1535)$ - resonance.

As an example on fig.5 the data obtained from the energy dependence $4\pi a_0(W)$ from paper [2] are shown.




Figure 5: The energy dependence of the real (a), imaginary (b) and phase φ (c) of the multipole amplitude E_{0+} from [2].

Upper and lower curves were calculated with error matrix of parameters W_R , $\sigma(W_R)$ Γ_R calculated with $b_\eta = 0.55$, $b_\pi = 0.35$ from table 2:

$$\begin{pmatrix} 1.54 & -1.23 & -4.71 \\ -1.23 & 2.16 & 1.84 \\ -4.71 & 1.84 & 33.20 \end{pmatrix}$$

Real and imaginary parts of E_{0+} transforms the regression coefficients containing E_{0+} into linear equations. Unfortunately the regression coefficients of polarization observables have significant errors and are at different energies [4]. This means that up to now there is no possibility to find the

reliable estimates of higher multipole. In order to go further and find reliable estimates of p -, d - and higher multipole it is necessary to have more precise and more detailed experimental data on polarization observables including angular distributions of recoiled protons polarization.

4 Conclusion

The partial wave analysis of observables of eta photoproduction on protons made in [4] confirms the dominating role of the s -wave in energy region from threshold up to about 2 GeV and demonstrates its resonance behaving corresponding to production of nucleon resonance $S_{11}(1535)$. The experimental data on differential cross sections don't show the presence of the partial waves with $l > 2$. In all cases the statistically reliable description of experimental data was achieved with keeping only three terms in expansion of differential cross section over Legendre polynomials. Contributions of p - and d -waves appear only through their interference with the resonance s -wave are small and the results of different experiments contradict with each other. Thus in the energy region up to 2 GeV the experimental data doesn't show appearance of large number of firmly established resonances [7].

The separation of s -wave contribution has allowed to find the energy dependence of amplitude E_{0+} and fundamental characteristics of $S_{11}(1535)$ resonance. Unfortunately the data of various experiments in this case are also incompatible.

References

- [1] B.Krusche et al., Phys.Rev.Lett. **74**, 3736 (1995); Phys.Rev.Lett. **75**, 3023 (1995)
- [2] F.Renard et al., Phys.Lett. **B528**, 215 (2002)
- [3] L.Soezueer et al., <http://gwdac.phys.gwu.edu>
- [4] E.V.Balandina et al., Yad.Phys., **65**, 1711 (2002); πN Newsletters **16**, 423 (2002); hep-ph 0111253
- [5] M.Dugger et al., PRL **89**, 222002 (2002)
- [6] E.V.Balandina et al., Preprint NPI MSU - 2002-12/696
- [7] Review of Partical Physics, <http://pdg.lbl.org>
- [8] H.Scheffe, The analysis of variance, N.-Y., John Wiley, 1958